

# Comment on “Quantum Suppression of Shot Noise in Atom-Size Metallic Contacts”

In a recent letter [1], van den Brom and van Ruitenbeek found a pronounced suppression of the shot noise in atom-size gold contacts with conductances near integer multiples of  $G_0 = 2e^2/h$ , revealing unambiguously the quantized nature of the electronic transport. However, the ad hoc model they introduced to describe the contribution of partially-open conductance channels to the shot noise is unable to fit either the maxima or minima of their shot noise data. Here we point out that a model of quantum-confined electrons with disorder [2] quantitatively reproduces the measurements of Ref. [1].

We model a nanocontact in a monovalent metal as a deformable constriction in an electron gas, with disorder included via randomly distributed delta-function potentials [2]. For convenience, the system is taken to be two-dimensional. The transmission probabilities  $T_n$  of the conducting channels are obtained via a modified recursive Green’s function algorithm [2]. The dimensionless shot noise  $s_I$  at zero temperature is [3]

$$s_I \equiv \frac{P_I}{2eI} = \frac{\sum_n T_n(1 - T_n)}{\sum_n T_n}, \quad (1)$$

where  $P_I$  is the shot noise spectral density and  $I$  is the time-average current. Plotting  $s_I$  versus the conductance  $G = G_0 \sum_n T_n$  eliminates the dependence on dimensionality for an ideal contact, provided no special symmetries are present.

Starting from the numerical data that were used to generate the conductance histogram in Ref. [2], we compute the mean and standard deviation of  $s_I$  and  $T_n$  as functions of  $G$ . The averages are taken over an ensemble of impurity configurations and contact shapes. The agreement of the experimental results for particular contacts and the calculated distribution of  $s_I$  shown in Fig. 1(b) is extremely good: 67% of the experimental points lie within one standard deviation of  $\langle s_I \rangle$  and 89% lie within two standard deviations [4]. It should be emphasized that no attempt has been made to fit the shot-noise data; the numerical data of Ref. [2], where the length of the contact and the strength of the disorder (mean-free path  $k_F \ell = 70$ ) were chosen to give qualitative agreement with experimental conductance histograms for gold [5], have simply been re-analyzed to calculate  $\langle s_I \rangle$ .

Contrary to the model of Ref. [1], the minima of  $\langle s_I \rangle$  do not occur at integer multiples of  $G_0$ , but are shifted to lower values, which correspond *not* to maxima of the conductance histogram, but rather to maxima of  $\langle T_n \rangle$ .

Fig. 1(c) shows that the number of partially-open channels increases in proportion to  $G$ . For comparison, the shot noise for a contact with only one partially-open channel, which sets a lower bound, is shown as a dashed curve in Fig. 1(b). The presence of several partially

open channels for  $G > G_0$  increases  $\langle s_I \rangle$  above this lower bound, leading to an apparent saturation at  $\langle s_I \rangle \approx 0.18$  for larger contacts. Neither the maxima nor the minima of the experimental data can be fit by the model of Ref. [1], which includes only two partially-open conductance channels.

The excellent agreement between the shot-noise data of Ref. [1] and our model calculation suggests that quantum transport in gold nanocontacts can be well described by a model which includes only two essential features, quantum confinement and coherent backscattering from imperfections in the contact.

J. B. acknowledges support from Swiss National Foundation PNR 36 “Nanosciences” grant # 4036-044033.

J. Bürki<sup>1,2,3</sup> and C. A. Stafford<sup>1</sup>

<sup>1</sup>University of Arizona, Tucson, Arizona 85721

<sup>2</sup>Université de Fribourg, 1700 Fribourg, Switzerland

<sup>3</sup>IRRMA, EPFL, 1015 Lausanne, Switzerland

Submitted to Phys. Rev. Lett. on 3 June 1999

PACS numbers: 72.70.+m, 72.15.Eb, 73.23.Ad, 73.40.Jn

- [1] H. E. van den Brom and J. M. van Ruitenbeek, Phys. Rev. Lett. **82**, 1526 (1999).
- [2] J. Bürki, C. A. Stafford, X. Zotos and D. Baeriswyl, cond-mat/9903006
- [3] M. Büttiker, Phys. Rev. Lett. **65**, 2901 (1990).
- [4] A finite temperature slightly lowers  $s_I$  for larger  $G$ , but does not significantly change the quality of the fit.
- [5] See e.g. J. L. Costa-Krämer, N. García, and H. Olin, Phys. Rev. B **55**, 12910 (1997).




FIG. 1. (a) Conductance histogram reproduced from Ref. [2]; (b) calculated mean shot noise  $\langle s_I \rangle$  (grey squares), together with experimental data from Ref. [1] (black circles); (c) mean transmission probabilities  $\langle T_n \rangle$ . The error bars indicate the standard deviations of the numerical results over the ensemble and the experimental errors, respectively.